


FIGURE 4: Sexual Assault HIV Exposure: Post-Exposure Prophylaxis (PEP) and Exposure Management When Reported Within 72 Hours
 Note: Regimens listed below are for individuals who weigh ≥ 40 kg; see Table 4 for PEP regimens for individuals who weigh < 40 kg.



Abbreviations: Ag/Ab, antigen/antibody; ART, antiretroviral therapy; HBV, hepatitis B virus; HCV, hepatitis C virus; PEP, post-exposure prophylaxis; STI, sexually transmitted infection.
Drug name abbreviations (brand name): 3TC, lamivudine (EpiVir); DTG, dolutegravir (Tivicay); FTC, emtricitabine (Emtriva); RAL, raltegravir (Isentress); TDF, tenofovir disoproxil fumarate (Viread).
Notes:
 a. Do not use fixed-dose combination tablet for patients who require dose adjustment for renal failure. Adjust dose of TDF/FTC or TDF/3TC for patients with creatinine clearance < 50 mL/min (see NYSDOH AI guideline Selecting an Initial ART Regimen > ARV Dose Adjustments for Hepatic or Renal Impairment).
 b. For HBV and HCV post-exposure management, see guideline sections Management of Potential Exposure to Hepatitis B Virus and Management of Potential Exposure to Hepatitis C Virus.