



## Laboratory Monitoring for Adverse Effects of ART

June 2021

**Table 1: Minimum Laboratory Monitoring Frequency With Initiation of or Change in ART for Patients <50 Years Old and Without Chronic Comorbidities [a] (Rating: A3)**

Laboratory Test	Year 1 of ART (initiation or change)			After 1 Year on ART Regimen	
	Baseline	3 Months	12 Months	Every 6 Months	Annual
Hepatic panel (AST, ALT, alkaline phosphates, total bilirubin)	All	All	All	—	All
Random blood glucose	All	All	All	—	—
Complete blood count [b]	All	With ZDV	With ZDV	With ZDV	—
eGFR [c]	All	All	With TAF or TDF	—	With TAF or TDF
Test for proteinuria (urinalysis or protein-to-creatinine ratio), glucosuria, serum phosphorus	With TAF or TDF	—	With TAF or TDF	—	With TAF or TDF

**Abbreviations:** ALT, alanine aminotransferase; ART, antiretroviral therapy; AST, aspartate aminotransferase; eGFR, estimated glomerular filtration rate; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate; ZDV, zidovudine.

**Notes:**

- a. More frequent monitoring may be required for patients >50 years old and patients with chronic comorbidities.
- b. See NYSDOH AI guideline [Primary Care for Adults With HIV](#).
- c. Patients with decreased eGFR at baseline or those taking concomitant nephrotoxic drugs may need more frequent monitoring of renal function (see guideline section [Screening for Organ-Specific Adverse Events > Nephrotoxicity](#) for more information).